

Table 7.7
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Recipient Age at Tx	Year of Transplant								
	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total									
Patients	155	218	301	302	374	344	419	342	292
Deaths	11	11	13	13	16	17	17	8	9
Rate	74.8	52.1	44.2	43.8	43.9	50.7	41.4	23.8	31.5
6-11 Years									
Patients	-	-	-	-	-	-	1	-	-
Deaths	-	-	-	-	-	-	0	-	-
Rate	-	-	-	-	-	-	*	-	-
18-34 Years									
Patients	32	40	65	50	73	52	86	62	50
Deaths	2	2	2	1	3	2	3	2	1
Rate	64.8	52.6	30.9	20.2	41.9	39.0	35.9	33.1	20.4
35-49 Years									
Patients	102	152	187	216	233	241	268	219	177
Deaths	6	7	9	11	7	13	13	5	5
Rate	61.4	47.5	49.4	52.1	30.7	55.4	49.6	23.3	28.7
50-64 Years									
Patients	21	26	49	35	67	51	64	60	64
Deaths	3	2	2	1	6	2	1	1	3
Rate	162.2	78.7	42.4	28.6	93.6	40.6	15.7	16.8	48.5
65+ Years									
Patients	-	-	-	1	1	-	-	1	1
Deaths	-	-	-	0	0	-	-	0	0
Rate	-	-	-	*	*	-	-	*	*

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Ethnicity/Race	Year of Transplant								
	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total									
Patients	155	218	301	302	374	344	419	342	292
Deaths	11	11	13	13	16	17	17	8	9
Rate	74.8	52.1	44.2	43.8	43.9	50.7	41.4	23.8	31.5
White									
Patients	147	199	274	254	319	299	367	282	256
Deaths	11	10	13	11	13	16	14	7	7
Rate	79.1	52.0	48.7	44.1	41.8	55.0	38.8	25.3	27.8
African American									
Patients	7	12	15	27	29	21	23	34	23
Deaths	0	1	0	2	2	0	3	1	2
Rate	*	86.7	0.0	76.3	71.2	0.0	144.6	29.7	93.3
Other/Multi-race									
Patients	-	4	1	3	1	4	1	2	1
Deaths	-	0	0	0	0	0	0	0	0
Rate	-	*	*	*	*	*	*	*	*
Asian									
Patients	-	-	-	2	3	4	3	2	1
Deaths	-	-	-	0	0	0	0	0	0
Rate	-	-	-	*	*	*	*	*	*
Hispanic/Latino									
Patients	1	3	11	16	22	16	25	22	11
Deaths	0	0	0	0	1	1	0	0	0
Rate	*	*	0.0	0.0	46.9	66.2	0.0	0.0	0.0

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Gender	Year of Transplant								
	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total									
Patients	155	218	301	302	374	344	419	342	292
Deaths	11	11	13	13	16	17	17	8	9
Rate	74.8	52.1	44.2	43.8	43.9	50.7	41.4	23.8	31.5
Female									
Patients	68	90	140	123	157	142	177	148	134
Deaths	3	3	5	4	9	6	9	5	3
Rate	45.2	34.2	36.2	33.1	58.9	43.3	52.4	34.6	22.8
Male									
Patients	87	128	161	179	217	202	242	194	158
Deaths	8	8	8	9	7	11	8	3	6
Rate	99.0	64.9	51.4	51.1	33.1	56.0	33.5	15.6	38.9

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Blood Type	Year of Transplant								
	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total									
Patients	155	218	301	302	374	344	419	342	292
Deaths	11	11	13	13	16	17	17	8	9
Rate	74.8	52.1	44.2	43.8	43.9	50.7	41.4	23.8	31.5
A									
Patients	80	94	126	132	165	134	163	134	132
Deaths	4	3	8	6	9	4	3	2	3
Rate	51.6	32.7	65.4	46.1	56.7	30.5	18.6	15.1	23.1
AB									
Patients	4	9	15	10	22	18	20	17	10
Deaths	1	0	0	1	0	3	1	0	0
Rate	*	*	0.0	100.1	0.0	186.4	51.9	0.0	0.0
B									
Patients	15	24	39	35	49	45	58	40	32
Deaths	1	0	2	0	2	1	3	2	2
Rate	71.1	0.0	53.7	0.0	41.2	22.3	52.7	51.4	65.4
O									
Patients	56	91	121	125	138	147	178	151	118
Deaths	5	8	3	6	5	9	10	4	4
Rate	95.3	92.7	25.1	49.4	36.9	63.0	57.9	27.1	34.6

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Residency	Year of Transplant								
	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total									
Patients	155	218	301	302	374	344	419	342	292
Deaths	11	11	13	13	16	17	17	8	9
Rate	74.8	52.1	44.2	43.8	43.9	50.7	41.4	23.8	31.5
U.S. Resident									
Patients	150	215	300	302	374	344	419	342	291
Deaths	11	10	13	13	16	17	17	8	9
Rate	77.4	47.9	44.4	43.8	43.9	50.7	41.4	23.8	31.6
Nonresident Alien									
Patients	1	-	-	-	-	-	-	-	1
Deaths	0	-	-	-	-	-	-	-	0
Rate	*	-	-	-	-	-	-	-	*
Unknown									
Patients	4	3	1	-	-	-	-	-	-
Deaths	0	1	0	-	-	-	-	-	-
Rate	*	*	*	-	-	-	-	-	-

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Hospitalized at Tx	Year of Transplant								
	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total									
Patients	155	218	301	302	374	344	419	342	292
Deaths	11	11	13	13	16	17	17	8	9
Rate	74.8	52.1	44.2	43.8	43.9	50.7	41.4	23.8	31.5
Not Hospitalized									
Patients	148	209	293	292	369	339	414	332	287
Deaths	11	11	13	13	15	17	17	7	8
Rate	78.5	54.5	45.5	45.4	41.6	51.5	41.9	21.4	28.4
Hospitalized									
Patients	4	6	5	8	5	5	4	8	5
Deaths	0	0	0	0	1	0	0	1	1
Rate	*	*	*	*	*	*	*	*	*
In Intensive Care									
Patients	2	2	3	2	-	-	1	2	-
Deaths	0	0	0	0	-	-	0	0	-
Rate	*	*	*	*	-	-	*	*	-
Unknown									
Patients	1	1	-	-	-	-	-	-	-
Deaths	0	0	-	-	-	-	-	-	-
Rate	*	*	-	-	-	-	-	-	-

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Donor Age	Year of Transplant								
	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total									
Patients	155	218	301	302	374	344	419	342	292
Deaths	11	11	13	13	16	17	17	8	9
Rate	74.8	52.1	44.2	43.8	43.9	50.7	41.4	23.8	31.5
<1 Year									
Patients	-	-	-	-	1	-	-	-	-
Deaths	-	-	-	-	0	-	-	-	-
Rate	-	-	-	-	*	-	-	-	-
1-5 Years									
Patients	-	1	-	-	-	-	-	-	-
Deaths	-	0	-	-	-	-	-	-	-
Rate	-	*	-	-	-	-	-	-	-
6-11 Years									
Patients	2	7	11	11	6	6	8	10	3
Deaths	0	1	1	0	1	1	1	0	0
Rate	*	*	94.4	0.0	*	*	*	0.0	*
12-17 Years									
Patients	25	44	59	59	72	75	94	74	52
Deaths	4	0	5	2	2	4	3	3	1
Rate	176.1	0.0	89.9	34.1	28.4	54.6	32.0	42.0	19.6
18-34 Years									
Patients	86	94	152	152	225	197	231	185	183
Deaths	4	3	3	9	10	10	8	5	6
Rate	48.2	32.2	19.9	60.8	45.5	52.3	35.5	27.5	33.5
35-49 Years									
Patients	36	64	63	69	60	62	83	70	52
Deaths	3	7	4	1	3	2	5	0	1
Rate	90.1	120.0	65.7	14.6	51.3	32.7	62.1	0.0	19.4
50-64 Years									
Patients	6	8	16	11	10	4	3	3	2
Deaths	0	0	0	1	0	0	0	0	1
Rate	*	*	0.0	97.9	0.0	*	*	*	*

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Level of HLA Mismatch (Total)		Year of Transplant								
		1998	1999	2000	2001	2002	2003	2004	2005	2006
Total	Patients	155	218	301	302	374	344	419	342	292
	Deaths	11	11	13	13	16	17	17	8	9
	Rate	74.8	52.1	44.2	43.8	43.9	50.7	41.4	23.8	31.5
A Locus Mismatch										
0	Patients	32	52	53	39	67	55	63	46	33
	Deaths	4	4	1	1	5	4	5	1	1
	Rate	138.9	80.8	19.0	25.8	79.6	75.9	83.4	22.2	30.9
1	Patients	91	105	137	147	183	168	195	163	134
	Deaths	6	4	5	7	6	6	3	2	4
	Rate	69.1	39.3	37.2	48.6	33.4	36.4	15.4	12.4	30.5
2	Patients	32	61	111	116	124	121	161	133	125
	Deaths	1	3	7	5	5	7	9	5	4
	Rate	31.8	50.3	65.5	43.9	41.0	59.4	57.6	38.7	32.7
Unknown	Patients	0	0	0	0	0	0	0	0	0
	Deaths	0	0	0	0	0	0	0	0	0
	Rate	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
B Locus Mismatch										
0	Patients	18	22	24	24	34	20	32	18	20
	Deaths	0	1	2	1	4	3	3	1	1
	Rate	0.0	47.5	88.2	43.1	131.2	165.7	100.3	58.5	51.6
1	Patients	75	95	124	124	130	144	151	125	102
	Deaths	5	6	3	4	5	6	5	3	3
	Rate	70.2	65.6	24.3	32.5	39.4	42.7	33.5	24.3	30.1
2	Patients	62	101	153	154	210	180	236	199	170
	Deaths	6	4	8	8	7	8	9	4	5
	Rate	103.7	40.6	54.0	53.2	33.8	45.3	39.0	20.4	30.0

(Continued)

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Mismatches are calculated without taking into account HLA from kidney donor.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Level of HLA Mismatch (Total)		Year of Transplant								
		1998	1999	2000	2001	2002	2003	2004	2005	2006
B Locus Mismatch										
Unknown										
	Patients	-	-	-	-	-	-	-	-	-
	Deaths	-	-	-	-	-	-	-	-	-
	Rate	-	-	-	-	-	-	-	-	-
DR Locus Mismatch										
0										
	Patients	27	31	44	38	42	41	49	38	32
	Deaths	1	2	1	0	3	2	2	2	2
	Rate	38.0	67.0	23.2	0.0	76.5	50.4	41.8	54.5	65.5
1										
	Patients	87	123	151	150	185	166	211	151	120
	Deaths	8	6	6	6	7	9	10	3	1
	Rate	98.1	50.7	40.6	41.0	38.7	55.5	48.5	20.2	8.4
2										
	Patients	41	64	106	113	146	137	159	153	140
	Deaths	2	3	6	6	6	6	5	3	6
	Rate	51.0	47.8	58.3	53.5	41.8	45.0	32.0	19.9	44.0
Unknown										
	Patients	-	-	-	1	1	-	-	-	-
	Deaths	-	-	-	1	0	-	-	-	-
	Rate	-	-	-	*	*	-	-	-	-
Total Mismatch										
0										
	Patients	2	6	3	6	12	7	11	5	6
	Deaths	0	0	0	0	3	1	2	1	1
	Rate	*	*	*	*	326.1	*	202.8	*	*
1										
	Patients	16	11	15	8	8	10	12	10	8
	Deaths	1	1	1	0	0	2	1	0	0
	Rate	65.4	99.3	71.0	*	*	222.2	90.4	0.0	*
2										
	Patients	19	31	41	35	31	36	43	20	16
	Deaths	2	2	2	1	1	1	1	1	1
	Rate	114.8	67.7	49.8	29.2	33.0	28.4	23.3	51.0	66.0

(Continued)

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Mismatches are calculated without taking into account HLA from kidney donor.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Level of HLA Mismatch (Total)		Year of Transplant								
		1998	1999	2000	2001	2002	2003	2004	2005	2006
Total Mismatch										
3										
	Patients	42	53	54	56	82	65	74	79	54
	Deaths	3	4	0	1	5	3	2	0	0
	Rate	75.8	78.5	0.0	18.1	63.8	47.5	27.4	0.0	0.0
4										
	Patients	43	57	83	78	100	99	104	77	67
	Deaths	2	2	3	5	2	4	2	1	2
	Rate	47.6	36.0	36.7	65.6	20.1	40.9	19.5	13.1	30.6
5										
	Patients	27	45	69	83	102	81	117	92	90
	Deaths	3	1	3	4	1	2	7	4	2
	Rate	121.3	22.7	44.3	48.9	9.8	24.8	61.1	44.9	22.6
6										
	Patients	6	15	36	35	38	46	58	59	51
	Deaths	0	1	4	1	4	4	2	1	3
	Rate	*	67.5	120.3	28.8	110.5	92.7	35.5	17.2	60.7
Unknown										
	Patients	-	-	-	1	1	-	-	-	-
	Deaths	-	-	-	1	0	-	-	-	-
	Rate	-	-	-	*	*	-	-	-	-

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Mismatches are calculated without taking into account HLA from kidney donor.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Cold Ischemic Time	Year of Transplant								
	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total									
Patients	155	218	301	302	374	344	419	342	292
Deaths	11	11	13	13	16	17	17	8	9
Rate	74.8	52.1	44.2	43.8	43.9	50.7	41.4	23.8	31.5
0-11 Hours									
Patients	26	38	65	74	90	82	129	119	114
Deaths	1	2	1	2	3	4	3	3	5
Rate	38.9	55.5	15.5	27.3	34.0	50.4	23.6	25.7	45.3
12-21 Hours									
Patients	85	110	142	130	165	149	166	138	122
Deaths	4	6	9	6	8	8	11	2	3
Rate	48.6	56.4	65.6	47.1	50.0	55.3	68.8	14.7	24.9
22-31 Hours									
Patients	23	36	28	23	38	20	32	27	16
Deaths	3	2	0	0	1	2	2	1	0
Rate	144.9	57.0	0.0	0.0	26.9	105.5	64.4	37.6	0.0
32-41 Hours									
Patients	1	3	-	-	1	-	2	1	4
Deaths	0	0	-	-	0	-	0	0	0
Rate	*	*	-	-	*	-	*	*	*
42+ Hours									
Patients	-	-	1	-	-	2	-	-	-
Deaths	-	-	0	-	-	0	-	-	-
Rate	-	-	*	-	-	*	-	-	-
Unknown									
Patients	20	31	65	75	80	91	90	57	36
Deaths	3	1	3	5	4	3	1	2	1
Rate	172.0	32.8	47.5	68.6	51.3	33.3	11.1	36.2	28.5

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Mismatches are calculated without taking into account HLA from kidney donor.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Transplant Center State		Year of Transplant								
		1998	1999	2000	2001	2002	2003	2004	2005	2006
Total	Patients	155	218	301	302	374	344	419	342	292
	Deaths	11	11	13	13	16	17	17	8	9
	Rate	74.8	52.1	44.2	43.8	43.9	50.7	41.4	23.8	31.5
AL	Patients	-	-	-	-	2	3	4	11	2
	Deaths	-	-	-	-	0	0	0	0	0
	Rate	-	-	-	-	*	*	*	0.0	*
AR	Patients	-	-	1	-	1	-	1	-	-
	Deaths	-	-	0	-	0	-	0	-	-
	Rate	-	-	*	-	*	-	*	-	-
AZ	Patients	-	1	1	3	8	2	12	15	7
	Deaths	-	0	1	0	0	0	1	0	0
	Rate	-	*	*	*	*	*	85.0	0.0	*
CA	Patients	2	9	22	11	26	28	25	16	20
	Deaths	0	0	0	0	0	6	0	0	0
	Rate	*	*	0.0	0.0	0.0	251.9	0.0	0.0	0.0
CO	Patients	-	-	1	1	3	2	1	1	1
	Deaths	-	-	0	0	0	0	0	0	0
	Rate	-	-	*	*	*	*	*	*	*
CT	Patients	4	4	3	9	4	5	2	2	3
	Deaths	0	0	0	1	0	0	0	0	0
	Rate	*	*	*	*	*	*	*	*	*
DC	Patients	6	2	1	4	5	4	5	4	2
	Deaths	0	1	0	1	0	0	0	0	1
	Rate	*	*	*	*	*	*	*	*	*
FL	Patients	1	2	9	18	15	11	10	2	6
	Deaths	1	0	1	0	1	0	1	0	0

(Continued)

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Mismatches are calculated without taking into account HLA from kidney donor.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Transplant Center State		Year of Transplant								
		1998	1999	2000	2001	2002	2003	2004	2005	2006
FL	Rate	*	*	*	0.0	67.0	0.0	101.5	*	*
GA	Patients	2	1	1	5	5	7	2	1	3
	Deaths	0	0	0	0	0	0	0	0	0
	Rate	*	*	*	*	*	*	*	*	*
HI	Patients	-	-	-	1	1	1	-	-	-
	Deaths	-	-	-	0	0	0	-	-	-
	Rate	-	-	-	*	*	*	-	-	-
IA	Patients	1	-	4	3	1	1	1	-	3
	Deaths	0	-	0	0	0	0	0	-	0
	Rate	*	-	*	*	*	*	*	-	*
IL	Patients	6	6	6	5	20	15	13	16	9
	Deaths	0	2	0	0	2	1	0	0	0
	Rate	*	*	*	*	109.1	69.5	0.0	0.0	*
IN	Patients	-	1	1	1	3	9	19	12	7
	Deaths	-	0	0	0	0	0	1	0	0
	Rate	-	*	*	*	*	*	54.9	0.0	*
KS	Patients	-	1	3	2	3	3	2	-	1
	Deaths	-	1	0	0	0	0	0	-	1
	Rate	-	*	*	*	*	*	*	-	*
KY	Patients	-	1	4	2	1	1	2	2	2
	Deaths	-	0	0	0	0	0	0	0	0
	Rate	-	*	*	*	*	*	*	*	*
LA	Patients	2	3	7	6	9	7	7	2	2
	Deaths	0	0	0	0	1	0	0	0	0
	Rate	*	*	*	*	*	*	*	*	*

(Continued)

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Mismatches are calculated without taking into account HLA from kidney donor.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Transplant Center State		Year of Transplant								
		1998	1999	2000	2001	2002	2003	2004	2005	2006
MA	Patients	2	3	20	11	11	17	25	19	16
	Deaths	0	0	0	0	0	1	0	1	0
	Rate	*	*	0.0	0.0	0.0	59.2	0.0	54.7	0.0
MD	Patients	16	34	23	19	19	13	16	17	13
	Deaths	2	1	2	3	1	0	1	1	0
	Rate	141.4	29.6	94.4	173.1	55.0	0.0	65.8	60.2	0.0
ME	Patients	-	-	-	-	2	3	6	1	4
	Deaths	-	-	-	-	0	1	0	0	0
	Rate	-	-	-	-	*	*	*	*	*
MI	Patients	6	9	9	8	11	4	14	15	15
	Deaths	0	1	0	0	0	0	3	0	2
	Rate	*	*	*	*	0.0	*	218.3	0.0	152.9
MN	Patients	62	72	79	96	65	55	52	50	39
	Deaths	5	3	3	4	3	6	2	0	0
	Rate	85.0	42.8	38.5	42.3	47.4	114.3	39.2	0.0	0.0
MO	Patients	-	-	-	-	-	1	4	-	6
	Deaths	-	-	-	-	-	0	0	-	0
	Rate	-	-	-	-	-	*	*	-	*
NC	Patients	2	6	7	6	14	18	10	11	3
	Deaths	0	0	1	0	0	0	1	0	0
	Rate	*	*	*	*	0.0	0.0	108.1	0.0	*
ND	Patients	-	-	-	2	10	7	6	2	3
	Deaths	-	-	-	0	0	0	0	0	0
	Rate	-	-	-	*	0.0	*	*	*	*
NE	Patients	4	3	4	4	7	3	7	1	3
	Deaths	0	0	0	0	0	0	0	0	0

(Continued)

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Mismatches are calculated without taking into account HLA from kidney donor.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Transplant Center State		Year of Transplant								
		1998	1999	2000	2001	2002	2003	2004	2005	2006
NE	Rate	*	*	*	*	*	*	*	*	*
NH	Patients	-	-	-	-	-	-	-	2	4
	Deaths	-	-	-	-	-	-	-	0	0
	Rate	-	-	-	-	-	-	-	*	*
NJ	Patients	1	-	-	4	17	19	22	7	4
	Deaths	0	-	-	0	1	0	2	1	1
	Rate	*	-	-	*	61.9	0.0	94.8	*	*
NV	Patients	1	-	-	-	-	-	-	-	-
	Deaths	0	-	-	-	-	-	-	-	-
	Rate	*	-	-	-	-	-	-	-	-
NY	Patients	4	3	16	11	15	13	18	11	12
	Deaths	0	0	1	1	0	1	1	0	0
	Rate	*	*	63.3	91.2	0.0	79.3	57.5	0.0	0.0
OH	Patients	5	7	21	18	18	16	26	28	27
	Deaths	0	0	1	1	2	0	2	2	0
	Rate	*	*	47.9	56.5	115.8	0.0	83.1	75.5	0.0
OK	Patients	-	-	-	-	1	3	2	-	1
	Deaths	-	-	-	-	0	0	0	-	0
	Rate	-	-	-	-	*	*	*	-	*
OR	Patients	-	1	-	1	3	4	1	-	5
	Deaths	-	0	-	0	0	0	0	-	0
	Rate	-	*	-	*	*	*	*	-	*
PA	Patients	5	13	17	8	20	17	27	18	14
	Deaths	0	1	2	0	3	1	0	2	0
	Rate	*	78.0	126.3	*	162.7	62.4	0.0	124.5	0.0

(Continued)

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Mismatches are calculated without taking into account HLA from kidney donor.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Transplant Center State		Year of Transplant								
		1998	1999	2000	2001	2002	2003	2004	2005	2006
RI	Patients	-	-	-	-	2	9	1	4	2
	Deaths	-	-	-	-	0	0	0	0	0
	Rate	-	-	-	-	*	*	*	*	*
SC	Patients	1	-	6	6	4	3	4	2	5
	Deaths	0	-	0	0	0	0	1	0	1
	Rate	*	-	*	*	*	*	*	*	*
SD	Patients	-	-	-	-	-	2	7	6	3
	Deaths	-	-	-	-	-	0	0	0	1
	Rate	-	-	-	-	-	*	*	*	*
TN	Patients	6	7	9	4	3	7	15	7	3
	Deaths	1	0	1	0	0	0	0	1	0
	Rate	*	*	*	*	*	*	0.0	*	*
TX	Patients	4	6	10	11	10	8	11	14	10
	Deaths	1	1	0	0	0	0	0	0	0
	Rate	*	*	0.0	0.0	0.0	*	0.0	0.0	0.0
UT	Patients	1	1	1	1	5	2	11	10	6
	Deaths	0	0	0	0	1	0	0	0	1
	Rate	*	*	*	*	*	*	0.0	0.0	*
VA	Patients	1	2	-	5	8	6	14	12	7
	Deaths	0	0	-	0	0	0	0	0	0
	Rate	*	*	-	*	*	*	0.0	0.0	*
VT	Patients	-	-	-	-	1	-	1	1	1
	Deaths	-	-	-	-	0	-	0	0	0
	Rate	-	-	-	-	*	-	*	*	*
WA	Patients	3	5	3	5	4	2	4	3	3
	Deaths	1	0	0	1	1	0	0	0	0

(Continued)

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Mismatches are calculated without taking into account HLA from kidney donor.

Table 7.7 (Continued)
Patients and Annual Death Rates Per 1,000 Patient-Years at Risk, 1998 to 2006
Rates for Recipients during First Year after Pancreas After Kidney (PAK) Transplantation

Transplant Center State		Year of Transplant								
		1998	1999	2000	2001	2002	2003	2004	2005	2006
WA	Rate	*	*	*	*	*	*	*	*	*
WI	Patients	7	15	12	11	17	13	9	17	15
	Deaths	0	0	0	1	0	0	1	0	1
	Rate	*	0.0	0.0	97.9	0.0	0.0	*	0.0	67.0

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

(-) = None in category.

(*) = For cohorts with fewer than 10 patients, rates are imprecise and are not reported.

Multi-organ transplants are excluded.

Includes patients transplanted during the year and having reported data one-year after transplant. Time at risk covers the period from transplant date until the date of death or one-year anniversary of the transplant. See Technical Notes for further details.

Mismatches are calculated without taking into account HLA from kidney donor.